

On (Super) Edge-Magic Total Labeling of Subdivision of $K_{1,3}$

Anak Agung Gede Ngurah,
Rinovia Simanjuntak and Edy Tri Baskoro

(Received August 31, 2006)

Abstract. Let G be a finite graph, with $V(G)$ and $E(G)$ the vertex-set and edge-set of G , respectively. An *edge-magic total labeling* is a one-to-one mapping f from $V(G) \cup E(G)$ onto $\{1, 2, 3, \dots, |V(G)| + |E(G)|\}$ such that there exists a constant c satisfying $f(u) + f(uv) + f(v) = c$, for each $uv \in E(G)$. Such a labeling is called a *super edge-magic total labeling* if all vertices of G receive all smallest labels. In this paper, we consider (super) edge-magic total labeling for subdivision of a star $K_{1,3}$.

AMS 2000 Mathematics Subject Classification. 05C78.

Key words and phrases. (Super) Edge-magic total labeling, subdivision of $K_{1,3}$.

§1. Introduction

All graphs considered here are finite and simple. The graph G has the vertex-set $V(G)$ and the edge-set $E(G)$.

Let $p = |V(G)|$ and $q = |E(G)|$. A bijection $f : V(G) \cup E(G) \rightarrow \{1, 2, 3, \dots, p + q\}$ is called an *edge-magic total labeling* of G if $f(x) + f(xy) + f(y)$ is a constant c (called the *magic constant* of f) for every edge xy of G . The graph that admits such a labeling is called an *edge-magic graph*. An edge-magic total labeling f is called a *super edge-magic total labeling* if $f(V(G)) = \{1, 2, 3, \dots, p\}$. A graph that admits a super edge-magic total labeling is called a *super edge-magic graph*. The edge-magic and super edge-magic concepts were first introduced by Kotzig and Rosa [7] and Enomoto, Lladó, Nakamigawa and Ringel [3], respectively.

Given a total labeling f , the *dual* labeling, which Kotzig and Rosa [7] called the complementary labeling, f' is defined as follows,

$$f'(x) = p + q + 1 - f(x) \text{ for every } x \in E(G) \cup V(G).$$

If f is an edge-magic total labeling with magic-constant c , then f' is an edge-magic total labeling with magic-constant $c' = 3(p + q + 1) - c$. Notice that this dual labeling does not preserve super edge-magic total labeling unless $G = \overline{K_n}$.

Another definition of a dual labeling was also introduced in [1]. By this definition the dual labeling preserves the property of super edge-magicness.

Lemma 1. [1] *If g is a super edge-magic total labeling of G with the magic constant c , then the function $g' : V(G) \cup E(G) \rightarrow \{1, 2, 3, \dots, p + q\}$ defined by*

$$g'(x) = \begin{cases} p + 1 - g(x), & \text{if } x \in V(G), \\ 2p + q + 1 - g(x), & \text{if } x \in E(G), \end{cases}$$

is also a super edge-magic total labeling of G with the magic constant $c' = 4p + q + 3 - c$.

The labeling g' defined in Lemma 1 is called a *dual super* labeling of g .

In the original papers about (super) edge-magic total labeling, Kotzig and Rosa [7], and Enomoto et al. [3] conjectured that every tree is edge-magic and every tree is super edge-magic, respectively. These conjectures have become very popular in the area of graph labeling. Some classes of tree have been proved to admit a (super) edge-magic labelings, such as paths, caterpillars [7], stars [4, 11], tree with at most 17 vertices [9], and path-like trees [2]. Additionally, Fukuchi [6] gives recursive formula for constructing super edge-magic trees. However, the conjectures are still remain open.

In this paper, we prove that a particular type of tree, namely a subdivision of a star $K_{1,3}$ is (super) edge-magic. These results provide more examples to support the correctness of the two conjectures on trees.

§2. The Results

For $m, n, k \geq 1$, let $T(m, n, k)$ be a graph obtained by inserting $m - 1$, $n - 1$, and $k - 1$ vertices to the first, second, and third edges, respectively, of a star $K_{1,3}$. Thus, the star $K_{1,3}$ can be written as $T(1, 1, 1)$. We define the vertex-set and the edge-set of graph $T(m, n, k)$ as follows.

$$V(T(m, n, k)) = \{w\} \cup \{x_i : 1 \leq i \leq m\} \cup \{y_i : 1 \leq i \leq n\} \cup \{z_i : 1 \leq i \leq k\},$$

and

$$E(T(m, n, k)) = \{wx_1, wy_1, wz_1\} \cup \{x_i x_{i+1} : 1 \leq i \leq m - 1\} \\ \cup \{y_i y_{i+1} : 1 \leq i \leq n - 1\} \cup \{z_i z_{i+1} : 1 \leq i \leq k - 1\}.$$

Clearly, a graph $T(m, n, k)$ has $m + n + k + 1$ vertices and $m + n + k$ edges. Among these vertices, one vertex has degree three, three vertices have degree one, and the remaining vertices have degree two. As an example, Figure 1 shows the graph $T(4, 5, 7)$.




Figure 1: A tree $T(4, 5, 7)$

Lu [12, 13] called the graph $T(m, n, k)$ as a three-path trees and proved that $T(m, n, k)$ is super edge-magic if n and k are odd, or $k = n + 1$, or $k = n + 2$. In this paper, we prove that $T(m, n, k)$ is also super edge-magic if $k = n + 3$, and $k = n + 4$.

In proving the main results, the following lemma will be frequently used.

Lemma 2. [4] *A graph G with p vertices and q edges is super edge-magic if and only if there exists a bijective function $f : V(G) \rightarrow \{1, 2, \dots, p\}$ such that the set $S = \{f(x) + f(y) : xy \in E(G)\}$ consists of q consecutive integers. In such a case, f extends to a super edge-magic total labeling of G with the magic constant $c = p + q + \min(S)$.*

Suppose $T(m, n, k)$ has an edge-magic total labeling with the magic constant c . Then tc , where $t = m + n + k$, cannot be smaller than the sum obtained by assigning the smallest labels to the vertex of degree 3, the $t - 3$ next smallest labels to the vertices of degree 2, and three next smallest labels to the vertices of degree 1; in other words

$$tc \geq 3 + 2 \sum_{i=2}^{t-2} i + \sum_{i=t-1}^{t+1} i + \sum_{i=t+2}^{2t+1} i.$$

An upper bound for tc is achieved by giving the the largest labels to the vertices of degree 3, and the $t - 3$ next largest labels to the vertices of degree 2, and 3 next largest labels to the rest of vertices; namely

$$tc \leq 3(2t + 1) + 2 \sum_{i=t+4}^{2t} i + \sum_{i=t+1}^{t+3} i + \sum_{i=1}^t i.$$

Thus, we have the following result.

Lemma 3. *If a $T(m, n, k)$ is an edge-magic graph, then magic constant c is in the following interval:*

$$\frac{1}{2t}(5t^2 + 3t + 6) \leq c \leq \frac{1}{2t}(7t^2 + 9t - 6).$$

By a similar argument, it is easy to verify that the following lemma holds.

Lemma 4. *If a $T(m, n, k)$ is a super edge-magic graph, then magic constant c is in the following interval:*

$$\frac{1}{2t}(5t^2 + 3t + 6) \leq c \leq \frac{1}{2t}(5t^2 + 11t - 6).$$

In the next two theorems, we will show that $T(m, n, k)$, for $k = n + 3$ and $k = n + 4$, is super edge-magic. First, we introduce two constants α and β used in the proposed labeling of graph $T(m, n, k)$ as follows:

$$\alpha = \begin{cases} 0, & \text{if } n \equiv 2 \pmod{4}, \\ 1, & \text{if } n \equiv 0, 1, 3 \pmod{4}, \end{cases}$$

and

$$\beta = \begin{cases} \lceil \frac{1}{2}(m-4) \rceil, & \text{if } n \equiv 0 \pmod{2}, \\ \lceil \frac{1}{2}(m-2) \rceil, & \text{if } n \equiv 1 \pmod{2}. \end{cases}$$

Theorem 1. *For all integers $m, n \geq 1$, $T(m, n, n + 3)$ is a super edge-magic graph.*

Proof. Consider the vertex labeling $f : V(T(m, n, n + 3)) \rightarrow \{1, 2, 3, \dots, m + 2n + 4\}$ defined as follows.

$$f(u) = \begin{cases} m + n + 3, & \text{if } u = w, \\ \lceil \frac{m}{2} \rceil - \frac{1}{2}(i-1), & \text{if } u = x_i \text{ for } i \equiv 1 \pmod{2}, \\ m + n + 3 - \frac{1}{2}i, & \text{if } u = x_i \text{ for } i \equiv 0 \pmod{2}, \\ \lceil \frac{m}{2} \rceil + 1 - \alpha + \frac{1}{2}(i+1), & \text{if } u = y_i \text{ for } i \equiv 1 \pmod{2}, \\ m + n + 3 + \frac{1}{2}i, & \text{if } u = y_i \text{ for } i \equiv 0 \pmod{2}. \end{cases}$$

For the remaining vertices, we consider the following four cases.

Case 1. $n \equiv 0 \pmod{4}$,

$$f(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + n + 1 - \frac{1}{2}(i-1), & \text{for } i \equiv 1 \pmod{4}, \\ \lceil \frac{m}{2} \rceil + n + 3 - \frac{1}{2}(i-1), & \text{for } i \equiv 3 \pmod{4}, \\ m + 2n + 4 - \frac{1}{2}i, & \text{for } i \equiv 2 \pmod{4}, i \neq n + 2, \\ m + 2n + 6 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{4}, \\ m + \frac{3}{2}n + 4, & \text{for } i = n + 2. \end{cases}$$

Case 2. $n \equiv 1 \pmod{4}$,

$$f(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + n + 1 - \frac{1}{2}(i-1), & \text{for } i \equiv 1 \pmod{4}, \\ \lceil \frac{m}{2} \rceil + n + 3 - \frac{1}{2}(i-1), & \text{for } i \equiv 3 \pmod{4}, \\ m + 2n + 4 - \frac{1}{2}i, & \text{for } i \equiv 2 \pmod{4}, \\ m + 2n + 6 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{4}. \end{cases}$$

Case 3. $n \equiv 2 \pmod{4}$,

$$f(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + 1, & \text{for } i = 1, \\ \lceil \frac{m}{2} \rceil + n + 3 - \frac{1}{2}(i - 1), & \text{for } i \equiv 1 \pmod{2}, i \neq 1, \\ m + 2n + 5 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{2}. \end{cases}$$

Case 4. $n \equiv 3 \pmod{4}$,

$$f(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + n + 1 - \frac{1}{2}(i - 1), & \text{for } i \equiv 1 \pmod{4}, i \neq n + 2, \\ \lceil \frac{m}{2} \rceil + n + 3 - \frac{1}{2}(i - 1), & \text{for } i \equiv 3 \pmod{4}, \\ m + 2n + 4 - \frac{1}{2}i, & \text{for } i \equiv 2 \pmod{4}, i \neq n - 1, \\ m + 2n + 6 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{4}, i \neq n + 1, \\ m + 5 + \frac{1}{2}(3n + 1), & \text{for } i = n - 1, \\ m + 3 + \frac{1}{2}(3n + 1), & \text{for } i = n + 1, \\ \lceil \frac{m}{2} \rceil + \lceil \frac{n}{2} \rceil + 1, & \text{for } i = n + 2, \\ m + 4 + \frac{1}{2}(3n + 1), & \text{for } i = n + 3. \end{cases}$$

Under the vertex labeling f , we have the following sums of labels of two adjacent vertices.

$$f(w) + f(x_1) = m + \lceil \frac{m}{2} \rceil + n + 3,$$

$$f(w) + f(y_1) = m + \lceil \frac{m}{2} \rceil + n + 5 - \alpha,$$


$$f(w) + f(z_1) = \begin{cases} m + \lceil \frac{m}{2} \rceil + n + 4, & \text{if } n \equiv 2 \pmod{4}, \\ m + \lceil \frac{m}{2} \rceil + 2n + 4, & \text{if } n \equiv 0, 1, 3 \pmod{4}, \end{cases}$$

$$\begin{aligned} & \{f(x_i) + f(x_{i+1}) : 1 \leq i \leq m - 1\} \\ &= \{\lceil \frac{m}{2} \rceil + n + 4, \lceil \frac{m}{2} \rceil + n + 5, \dots, m + \lceil \frac{m}{2} \rceil + n + 2\}, \\ & \{f(y_i) + f(y_{i+1}) : 1 \leq i \leq n - 1\} \\ &= \{m + \lceil \frac{m}{2} \rceil + n + 6 - \alpha, m + \lceil \frac{m}{2} \rceil + n + 7 - \alpha, \dots, m + \lceil \frac{m}{2} \rceil + 2n + 4 - \alpha\}, \\ & \{f(z_i) + f(z_{i+1}) : 1 \leq i \leq n + 2\} \\ &= \{m + \lceil \frac{m}{2} \rceil + 2n + 5, m + \lceil \frac{m}{2} \rceil + 2n + 6, \dots, m + \lceil \frac{m}{2} \rceil + 3n + 6\}. \end{aligned}$$

Thus, the set $S = \{f(v) + f(w) : vw \in E(T(m, n, n + 3))\}$ consists of consecutive integers with $\max(S) = m + \lceil \frac{m}{2} \rceil + 3n + 6$. By Lemma 2, f extends to a super edge-magic total labeling of $T(m, n, n + 3)$ with magic constant $c = 2m + \lceil \frac{m}{2} \rceil + 5n + 11$. \square

Figure 2 shows the vertex labeling of a super edge-magic tree $T(4, 6, 9)$.

Theorem 2. *For all integers $m, n \geq 1$, $T(m, n, n + 4)$ is a super edge-magic graph.*

Figure 2: A super edge-magic tree $T(4, 6, 9)$

Proof. Label the vertices of $T(m, n, n + 4)$ in the following way. We consider 2 cases, where n is even and odd.

Case 1. $n \equiv 0 \pmod{2}$,

$$g(u) = f(u), \quad \text{for } u = w, x_i^{\epsilon} s, \text{ and } y_i^{\epsilon} s,$$

where f is the vertex labeling in the proof of Theorem 1 with $\alpha = 1$.

Subcase 1.1. $n \equiv 0 \pmod{4}$,

$$g(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + n + 1 - \frac{1}{2}(i - 1), & \text{for } i \equiv 1 \pmod{4}, \\ \lceil \frac{m}{2} \rceil + n + 3 - \frac{1}{2}(i - 1), & \text{for } i \equiv 3 \pmod{4}, \\ m + 2n + 5 - \frac{1}{2}i, & \text{for } i \equiv 2 \pmod{4}, \\ m + 2n + 7 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{4}. \end{cases}$$

Subcase 1.2. $n \equiv 2 \pmod{4}$,

$$g(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + n + 1 - \frac{1}{2}(i - 1), & \text{for } i \equiv 1 \pmod{4}, i \neq n + 3, \\ \lceil \frac{m}{2} \rceil + n + 3 - \frac{1}{2}(i - 1), & \text{for } i \equiv 3 \pmod{4}, \\ m + 2n + 5 - \frac{1}{2}i, & \text{for } i \equiv 2 \pmod{4}, i \neq n, n + 4, \\ m + 2n + 7 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{4}, i \neq n + 2, \\ m + 6 + \frac{3}{2}n, & \text{for } i = n, \\ m + 4 + \frac{3}{2}n, & \text{for } i = n + 2, \\ \lceil \frac{m+n}{2} \rceil + 1, & \text{for } i = n + 3, \\ m + 5 + \frac{3}{2}n, & \text{for } i = n + 4. \end{cases}$$

Case 2. $n \equiv 1 \pmod{2}$,

$$g(u) = \begin{cases} m + n + 4, & \text{if } u = w, \\ \lceil \frac{m}{2} \rceil - \frac{1}{2}(i - 1), & \text{if } u = x_i \text{ for } i \equiv 1 \pmod{2}, \\ m + n + 4 - \frac{1}{2}i, & \text{if } u = x_i \text{ for } i \equiv 0 \pmod{2}, \\ \lceil \frac{m}{2} \rceil + 2 + \frac{1}{2}(i - 1), & \text{if } u = y_i \text{ for } i \equiv 1 \pmod{2}, \\ m + n + 4 + \frac{1}{2}i, & \text{if } u = y_i \text{ for } i \equiv 0 \pmod{2}. \end{cases}$$

$$g(z_i) = \begin{cases} \lceil \frac{m}{2} \rceil + 1, & \text{for } i = 1, \\ \lceil \frac{m}{2} \rceil + n + 4 - \frac{1}{2}(i - 1), & \text{for } i \equiv 1 \pmod{2}, i \neq 1, \\ m + 2n + 6 - \frac{1}{2}i, & \text{for } i \equiv 0 \pmod{2}. \end{cases}$$

It is a routine procedure to verify that g is a vertex labeling of $T(m, n, n+4)$. Under the vertex labeling g , for each case of n , we can count the sums of labels of two adjacent vertices as follows.

Case 1. $n \equiv 0 \pmod{2}$,

$$g(w) + g(x_1) = m + \left\lceil \frac{m}{2} \right\rceil + n + 3,$$

$$g(w) + g(y_1) = m + \left\lceil \frac{m}{2} \right\rceil + n + 4,$$

$$g(w) + g(z_1) = m + \left\lceil \frac{m}{2} \right\rceil + 2n + 4,$$

$$\begin{aligned} &\{g(x_i) + g(x_{i+1}) : 1 \leq i \leq m-1\} \\ &= \{\left\lceil \frac{m}{2} \right\rceil + n + 4, \left\lceil \frac{m}{2} \right\rceil + n + 5, \dots, m + \left\lceil \frac{m}{2} \right\rceil + n + 2\}, \\ &\{g(y_i) + g(y_{i+1}) : 1 \leq i \leq n-1\} \\ &= \{m + \left\lceil \frac{m}{2} \right\rceil + n + 5, m + \left\lceil \frac{m}{2} \right\rceil + n + 6, \dots, m + \left\lceil \frac{m}{2} \right\rceil + 2n + 3\}, \\ &\{g(z_i) + g(z_{i+1}) : 1 \leq i \leq n+3\} \\ &= \{m + \left\lceil \frac{m}{2} \right\rceil + 2n + 5, m + \left\lceil \frac{m}{2} \right\rceil + 2n + 6, \dots, m + \left\lceil \frac{m}{2} \right\rceil + 3n + 7\}. \end{aligned}$$

Case 2. $n \equiv 1 \pmod{2}$,

$$g(w) + g(x_1) = m + \left\lceil \frac{m}{2} \right\rceil + n + 4,$$

$$g(w) + g(y_1) = m + \left\lceil \frac{m}{2} \right\rceil + n + 6,$$


$$g(w) + g(z_1) = m + \left\lceil \frac{m}{2} \right\rceil + n + 5,$$

$$\begin{aligned} &\{g(x_i) + g(x_{i+1}) : 1 \leq i \leq m-1\} \\ &= \{\left\lceil \frac{m}{2} \right\rceil + n + 5, \left\lceil \frac{m}{2} \right\rceil + n + 6, \dots, m + \left\lceil \frac{m}{2} \right\rceil + n + 3\}, \\ &\{g(y_i) + g(y_{i+1}) : 1 \leq i \leq n-1\} \\ &= \{m + \left\lceil \frac{m}{2} \right\rceil + n + 7, m + \left\lceil \frac{m}{2} \right\rceil + n + 8, \dots, m + \left\lceil \frac{m}{2} \right\rceil + 2n + 5\}, \\ &\{g(z_i) + g(z_{i+1}) : 1 \leq i \leq n+3\} \\ &= \{m + \left\lceil \frac{m}{2} \right\rceil + 2n + 6, m + \left\lceil \frac{m}{2} \right\rceil + 2n + 7, \dots, m + \left\lceil \frac{m}{2} \right\rceil + 3n + 8\}. \end{aligned}$$

Hence, the set $S = \{g(v) + g(w) : vw \in E(T(m, n, n+4))\}$ is a set of consecutive integers with $\max(S) = m + 3n + 9 + \beta$. By Lemma 2, g extends to a super edge-magic total labeling of $T(m, n, n+4)$ with magic constant $c = 2m + 5n + 15 + \beta$. \square

Figure 3 shows the vertex labeling of a super edge-magic tree $T(3, 6, 10)$.

By the dual super property (Lemma 1), $T(m, n, n+3)$ and $T(m, n, n+4)$ also have a super edge-magic total labeling with magic constant as in the following corollaries.

Figure 3: A super edge-magic tree $T(3, 6, 10)$

Corollary 1. For all integers $m, n \geq 1$, $T(m, n, n+3)$ has a super edge-magic total labeling with magic constants $c = 3m - \lceil \frac{m}{2} \rceil + 5n + 11$. \square

Corollary 2. For all integers $m, n \geq 1$, $T(m, n, n+4)$ has a super edge-magic total labeling with magic constants $c = 3m + 5n + 12 - \beta$. \square

Additionally, by applying the duality property to Theorems 1 and 2, and Corollaries 1 and 2, we have the following results.

Corollary 3. For all integers $m, n \geq 1$, $T(m, n, n+3)$ has edge-magic total labelings with magic constants $c = 4m - \lceil \frac{m}{2} \rceil + 7n + 13$ and $c = 3m + \lceil \frac{m}{2} \rceil + 7n + 13$. \square

Corollary 4. For all integers $m, n \geq 1$, $T(m, n, n+4)$ has edge-magic total labelings with magic constants $c = 4m + 7n + 15 - \beta$ and $c = 3m + 7n + 18 + \beta$. \square

We can also construct an edge-magic total labeling of $T(m, n, n+3)$ and $T(m, n, n+4)$ for which all the odd labels are on the vertices, as follows.

Theorem 3. For all integers $m, n \geq 1$, $T(m, n, n+3)$ has an edge-magic total labeling with all vertices receive odd labels. This labeling has magic constant $c = 2m + 2\lceil \frac{m}{2} \rceil + 6n + 12$ and the dual has magic constant $c = 4m - 2\lceil \frac{m}{2} \rceil + 6n + 12$.

Proof. Define a labeling h of $T(m, n, n+3)$ as follows.

$$h(v) = 2f(v) - 1, \text{ for all } v \in V(T(m, n, n+3)),$$

where f is the vertex labeling in the proof of Theorem 1. It is not difficult to verify that all vertices receive odd labels, and

$$S = \{h(u) + h(v) : uv \in E(T(m, n, n+3))\}$$

forms an arithmetic progression with initial term $2n + 2\lceil \frac{m}{2} \rceil + 6$ having common difference 2. If we define

$$h(uv) = 2m + 2\lceil \frac{m}{2} \rceil + 6n + 12 - h(u) - h(v),$$

then h is an edge-magic total labeling of $T(m, n, n+3)$ with magic constant $c = 2m + 2\lceil \frac{m}{2} \rceil + 6n + 12$. By the duality property, it also has an edge-magic total labeling with magic constant $c = 4m - 2\lceil \frac{m}{2} \rceil + 6n + 12$. \square

A similar result for $T(m, n, n+4)$ can be stated in the next theorem.

Theorem 4. *For all integers $m, n \geq 1$, $T(m, n, n+4)$ has an edge-magic total labeling with all vertices receive odd labels. This labeling has magic constant $c = 2m + 6n + 18 + 2\beta$ and the dual has magic constant $c = 4m + 6n + 12 - 2\beta$, where β is a constant as defined before.*

We have proved the super edge-magicness of $T(m, n, k)$ only for $k = n+3$ and $k = n+4$ (not for any value of k). Additionally, we proved that $T(m, n, n+3)$ and $T(m, n, n+4)$ are (super) edge-magic for several values of magic constants c but not for all possible values of c . So, we have the following open problems.

Open problem 1. *Find a (super) edge-magic total labeling of $T(m, n, k)$ for any remaining values of m, n and k .*

Open problem 2. *Find a (super) edge-magic total labeling of $T(m, n, n+3)$ and $T(m, n, n+4)$ for other values of magic constants c .*

Acknowledgement

The authors would like to thank the referee for his/her valuable comments.

References

- [1] E. T. Baskoro, I. W. Sudarsana and Y. M. Cholily, How to construct new super edge-magic graphs from some old ones, *J. Indones. Math. Soc. (MIHMI)*, **11**:2 (2005), 155–162.
- [2] M. Baca, Y. Lin and F. A. Muntaner-Batle, Super edge-antimagic labelings of the path-like trees, *Utilitas Math.*, to appear.
- [3] H. Enomoto, A. Lladó, T. Nakamigawa, and G. Ringel, Super edge magic graphs, *SUT J. Math.*, **34** (1998), 105–109.
- [4] R. M. Figueroa-Centeno, R. Ichishima and F. A. Muntaner-Batle, The place of super edge-magic labelings among other classes of labelings, *Discrete Math.*, **231** (2001), 153–168.
- [5] R. M. Figueroa-Centeno, R. Ichishima and F. A. Muntaner-Batle, On edge-magic labelings of certain disjoint union graphs, *Australas. J. Combin.*, **32** (2005), 225–242.

- [6] Y. Fukuchi, A recursive theorems for super edge-magic labelings of trees, *SUT J. Math.*, **36**:2 (2000), 279–285.
- [7] A. Kotzig and A. Rosa, Magic valuation of finite graphs, *Canad. Math. Bull.*, Vol. **13**:4, (1970), 451–461.
- [8] A. Kotzig and A. Rosa, Magic valuation of complete graphs, *Publications du Centre de Recherches mathématiques Université de Montréal*, **175** (1972).
- [9] S. M. Lee, and Q. X. Shan, All trees with at most 17 vertices are super edge-magic, *16th MCCCC Conf.*, Carbondale, University Southern Illinois, Nov. 2002.
- [10] Slamin, M. Băca, Y. Lin, M. Miller and R. Simanjuntak, Edge magic total labeling of wheels, fans and friendship graphs, *Bull. Inst. Combin. Appl.*, **35** (2002), 89–98.
- [11] W. D. Wallis, E. T. Baskoro, M. Miller and Slamin, Edge-magic total labelings, *Austral. J. Combin.*, **22** (2000), 177–190.
- [12] Yong-Ji Lu, A proof of three-path trees $P(m, n, t)$ being edge-magic, *College Mathematics*, **17**:2 (2001), 41–44.
- [13] Yong-Ji Lu, A proof of three-path trees $P(m, n, t)$ being edge-magic (II), *College Mathematics*, **20**:3 (2004), 51–53.

Anak Agung Gede Ngurah

Department of Civil Engineering, Universitas Merdeka Malang

Jl. Taman Agung No.1 Malang, Indonesia

and

Combinatorial Mathematics Research Group, Faculty of Mathematics and Natural Sciences,

Institut Teknologi Bandung

Jalan Ganesa 10 Bandung, Indonesia

E-mail: s304agung@dns.math.itb.ac.id

Rinovia Simanjuntak and Edy Tri Baskoro

Combinatorial Mathematics Research Group, Faculty of Mathematics and Natural Sciences,

Institut Teknologi Bandung

Jalan Ganesa 10 Bandung, Indonesia

E-mail: {rino, ebaskoro}@dns.math.itb.ac.id